





<p><b>Student's obligation</b></p>	<p>Attending the lecture is a fundamental part of the course. You are responsible for material presented in the lecture whether or not it is discussed in the textbook. You should expect questions on the exams to test your understanding of concepts discussed in the lecture and in the homework assignments.</p> <p>It can be very helpful to study with a group. This type of cooperative learning is encouraged; however, be sure that you have a thorough understanding of the concepts besides the mathematical steps used to solve a problem. You must be able to work through the problems on your own.</p>			
<p><b>Required Learning Materials</b></p>	<p>Data Show, Handout lecture notes and white board notes.</p>			
<p><b>Evaluation</b></p>	<p><b>Task</b></p>	<p><b>Weight (Marks)</b></p>	<p><b>Due Week</b></p>	<p><b>Relevant Learning Outcome</b></p>
<p>Paper Review</p>				
<p style="writing-mode: vertical-rl; transform: rotate(180deg);">Assignments</p>		<p>Homework</p>	<p>14%</p>	
		<p>Class Activity</p>	<p>2%</p>	
		<p>Report</p>	<p>8%</p>	
		<p>Seminar</p>	<p>8%</p>	
		<p>Essay</p>		
		<p>Project</p>	<p>8%</p>	
<p>Quiz</p>		<p>4%</p>		
<p>Lab.</p>				
<p>Midterm Exam</p>		<p>16%</p>		
<p>Final Exam</p>		<p>40%</p>		
<p>Total</p>		<p>100%</p>		
<p><b>Specific learning outcome:</b></p>	<p>Upon completion of the course, the student will be able to:</p> <ol style="list-style-type: none"> <li>1. Find the anti-derivative of elementary polynomials, exponential, logarithmic and trigonometric functions.</li> <li>2. Use the Fundamental Theorem of Calculus to evaluate definite integrals.</li> <li>3. Definite integrals to find areas of planar regions.</li> <li>4. Interpret the definite integral geometrically as the area under a curve.</li> <li>5. Construct a definite integral as the limit of a Riemann sum.</li> </ol>			



<b>Course References:</b>	<p>1. Stewart, J. (2016). <i>Single variable calculus: Early transcendentals</i> (Eight edition.). Boston, MA, USA: Cengage Learning. (Major).</p> <p>2. Lecture Notes. (Minor)</p>	
<b>Course topics (Theory)</b>	<b>Week</b>	<b>Learning Outcome</b>
Integration, Indefinite Integrals and definite integrals.	1	
Integration of Trigonometric Function and Exponential Function	2	
Substitution Rule	3	
Integration by Parts	4	
Integration of Rational Functions by Partial Fractions	5	
Numerical Integration: Trapezoidal Rule and Simpson's Rule.	6 & 7	
Area, Area Between Curves and Average Value of a Continuous Function Revisited	8	
Volume, Volume of Pyramid, Volume of a Wedge, A Solid of Revolution, Volume of Sphere,	9	
Differential Equations	10 and 11	
Mid Term Exam	12	
Final Exam	12	

## Questions Example Design

Q1:

$$\text{Find } \int \cos (7\theta + 3) d\theta.$$

**Solution:**

$$\begin{aligned} \int \cos (7\theta + 3) d\theta &= \frac{1}{7} \int \cos (7\theta + 3) \cdot 7 d\theta \\ &= \frac{1}{7} \int \cos u du \\ &= \frac{1}{7} \sin u + C \\ &= \frac{1}{7} \sin (7\theta + 3) + C \end{aligned}$$

Q2:

Use partial fractions to evaluate

$$\int \frac{dx}{x(x^2 + 1)^2}.$$

**Solution** The form of the partial fraction decomposition is

$$\frac{1}{x(x^2 + 1)^2} = \frac{A}{x} + \frac{Bx + C}{x^2 + 1} + \frac{Dx + E}{(x^2 + 1)^2}$$

Multiplying by  $x(x^2 + 1)^2$ , we have

$$\begin{aligned} 1 &= A(x^2 + 1)^2 + (Bx + C)x(x^2 + 1) + (Dx + E)x \\ &= A(x^4 + 2x^2 + 1) + B(x^4 + x^2) + C(x^3 + x) + Dx^2 + Ex \\ &= (A + B)x^4 + Cx^3 + (2A + B + D)x^2 + (C + E)x + A \end{aligned}$$

If we equate coefficients, we get the system

$$A + B = 0, \quad C = 0, \quad 2A + B + D = 0, \quad C + E = 0, \quad A = 1.$$

Solving this system gives  $A = 1, B = -1, C = 0, D = -1$ , and  $E = 0$ . Thus,

$$\begin{aligned} \int \frac{dx}{x(x^2 + 1)^2} &= \int \left[ \frac{1}{x} + \frac{-x}{x^2 + 1} + \frac{-x}{(x^2 + 1)^2} \right] dx \\ &= \int \frac{dx}{x} - \int \frac{x dx}{x^2 + 1} - \int \frac{x dx}{(x^2 + 1)^2} \\ &= \int \frac{dx}{x} - \frac{1}{2} \int \frac{du}{u} - \frac{1}{2} \int \frac{du}{u^2} \quad \begin{array}{l} u = x^2 + 1, \\ du = 2x dx \end{array} \\ &= \ln |x| - \frac{1}{2} \ln |u| + \frac{1}{2u} + K \\ &= \ln |x| - \frac{1}{2} \ln (x^2 + 1) + \frac{1}{2(x^2 + 1)} + K \\ &= \ln \frac{|x|}{\sqrt{x^2 + 1}} + \frac{1}{2(x^2 + 1)} + K. \end{aligned}$$

**Q3: Find the integrals of the following:**

- $\int_1^4 \left( \frac{3}{2} \sqrt{x} - \frac{4}{x^2} \right) dx$
- $\int x \cos x dx.$

**Solution: 1)**

$$\int_1^4 \left( \frac{3}{2} \sqrt{x} - \frac{4}{x^2} \right) dx = \left[ x^{3/2} + \frac{4}{x} \right]_1^4 \qquad \frac{d}{dx} \left( x^{3/2} + \frac{4}{x} \right) = \frac{3}{2} x^{1/2} - \frac{4}{x^2}$$

$$= \left[ (4)^{3/2} + \frac{4}{4} \right] - \left[ (1)^{3/2} + \frac{4}{1} \right]$$

$$= [8 + 1] - [5] = 4$$

2)

**Solution** We use the formula  $\int u dv = uv - \int v du$  with

$$u = x, \quad dv = \cos x dx,$$

$$du = dx, \quad v = \sin x. \qquad \text{Simplest antiderivative of } \cos x$$

Then


$$\int x \cos x dx = x \sin x - \int \sin x dx = x \sin x + \cos x + C.$$

**Q4:**

Approximate the area under the curve  $y = \frac{1}{x}$  between  $x = 1$  and  $x = 5$  using the Trapezoidal Rule with  $n = 4$  subintervals.



*Solution.*



We write the Trapezoidal Rule formula for  $n = 4$  subintervals:

$$T_4 = \frac{\Delta x}{2} [f(x_0) + 2f(x_1) + 2f(x_2) + 2f(x_3) + f(x_4)].$$

The function has the following values at the points  $x_i$  :

$$f(x_0) = f(1) = \frac{1}{1} = 1;$$

$$f(x_1) = f(2) = \frac{1}{2};$$

$$f(x_2) = f(3) = \frac{1}{3};$$

$$f(x_3) = f(4) = \frac{1}{4};$$

$$f(x_4) = f(5) = \frac{1}{5}.$$

$$f(x_4) = f(5) = \frac{1}{5}.$$

Since  $\Delta x = 1$ , we obtain

$$\begin{aligned} A \approx T_4 &= \frac{1}{2} \left[ 1 + 2 \cdot \frac{1}{2} + 2 \cdot \frac{1}{3} + 2 \cdot \frac{1}{4} + \frac{1}{5} \right] = \frac{1}{2} \left[ 1 + 1 + \frac{2}{3} + \frac{1}{2} + \frac{1}{5} \right] \\ &= \frac{1}{2} \cdot \frac{30 + 30 + 20 + 15 + 8}{30} = \frac{1}{2} \cdot \frac{101}{30} = \frac{101}{60} \end{aligned}$$

**Extra notes:**

**External Evaluator**

Approved

Dr. Saad Khalis Essa

